



# Migrate™

GOOSE & BIRD REPELLENT

For Agricultural Use: To be used on Agricultural Commodities: Berries, Apples, Stone Fruits, Cereal Grains, and Sunflowers and Non-Agricultural Uses including Turf, Hydroseeding, Golfcourse, Food Stuffs, Wares & Traps, and Sport Repelling

ACTIVE INGREDIENT: N-ethylmalethylmalethylamine ..... 14.9%  
OTHER INGREDIENTS ..... 85.1%  
TOTAL ..... 100.0%  
This product contains 1.23 pounds/0.31 pounds/0.15 pounds of N-ethylmalethylmalethylamine per gallon/quarter pint/pint of product.

EPA REG. NO. 58035-9      EPA EST. NO. 071638-08-001  
EPA REG. NO. 073273-01-001

KEEP OUT OF REACH OF CHILDREN  
CAUTION

PRELIMINARY	
For Eyes	• If eyes open and close slowly and gently with water for 15-30 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
For Skin or Clothing	• Take off contaminated clothing. • Wash skin immediately with plenty of water for 15-30 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300. From agency or retail store, obtain safety data sheet.	

Manufactured by: Chemagro, Inc. #70 Box 2404 • Gaston, NC 28053

Toll Free 866-632-4737 • 260-447-4190 • FAX 260-447-7827

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS  
CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS  
For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to terrestrial areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or residue.

PERSONAL PROTECTIVE EQUIPMENT (PPE):  
Applicators and other handlers must wear:  
• Long-sleeved shirt and long pants  
• W neoprene gloves  
• Shoes plus socks.  
Follow manufacturer's instructions for cleaning/maintaining PPE. Keep and wash PPE separately from other laundry.

User Safety Recommendations Users Should:
• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE  
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact others or other persons, either directly or through drift, only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this labeling about PPE, restricted-entry intervals, and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS). Do not enter or allow workers into treated areas during the restricted-entry interval (REI) of 8 hours.

Specimen Label

NON-AGRICULTURAL USE REQUIREMENTS
The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep unprotected persons out of treated areas until sprays have dried.

AGRICULTURAL CROPS

USE RESTRICTIONS:  
Use this product to limit damage caused by birds on opening crops. This product must be applied using appropriate spray equipment. Do not apply when crop is wet or rain is expected. Wet foliage remains until fruit is susceptible to bird damage. Do not apply through any type of irrigation system.

COMPATIBILITY WITH OTHER PESTICIDES:  
Compatibility with other pesticides has not been tested. If adding other pesticides or products, test for compatibility first or ask your representative for advice. Do not mix with dry formulations such as wettable powder formulations which may result in residue on the crop.

DILUTION DIRECTIONS:  
Shake or stir RejeX-4M concentrate well prior to diluting. For better coverage, add a surfactant (such as a non-ionic, organofunctional silicone surfactant, or sticker/spreader), as directed on the label. Mix product outside or in a ventilated area. For ground application, mix RejeX-4M concentrate with water at a ratio of 1 part product to 10-20 parts water. For aerial application, mix RejeX-4M concentrate with water at a ratio of 1 part product to 1-3 parts water.

APPLICATION DIRECTIONS:  
• Apply spray mixture so that crop receives thorough coverage.  
• Repeat treatment if significant bird damage begins to occur.  
• This product may be applied up to the day of harvest.

USE SITE	APPLICATION DIRECTIONS	NOTES
Apples	Apply by ground application at a rate of 2.5 - 5 gallons of RejeX-4M concentrate (0.1-0.2 lbs a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:10 or 1:20 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	
Stone Fruits Apricots Cherries Peaches Plums	Apply by ground application at a rate of 2.5 - 5 gallons of RejeX-4M concentrate (0.1-0.2 lbs a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:10 or 1:20 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	California only: Mix concentrate and at least application mixture coverable over most of crop and not prepare spray solutions using more than 50 gallons of concentrate per day. Harvesters must wear gloves and long-sleeved shirts.
Blueberries	Apply by ground application at a rate of 2.5 - 5 gallons of RejeX-4M concentrate (0.1-0.2 lbs a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:10 or 1:20 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	California only: Mix concentrate and at least application mixture coverable over most of crop and not prepare spray solutions using more than 50 gallons of concentrate per day. Harvesters must wear gloves and long-sleeved shirts.
Grapes (table)	Apply by ground application at a rate of 2.5 - 5 gallons of RejeX-4M concentrate (0.1-0.2 lbs a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:10 or 1:20 dilution with water and apply at rates of 27.5-55 gallons or 52.5-100 gallons, respectively, of spray mixture per acre.	California only: Mix concentrate and at least application mixture coverable over most of crop and not prepare spray solutions using more than 50 gallons of concentrate per day. Harvesters must wear gloves and long-sleeved shirts. New York only: Aerial application is prohibited. This product is not to be used on grapes within Sullivan County.
Cereal Grains Corn (including sweet and pop) Millet Oats Sorghum	Apply by aerial application at a rate of 2.5 - 5 gallons of RejeX-4M concentrate (0.1-0.2 lbs a.i.) per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:10 dilution with water and apply at rates of 10-20 gallons of spray mixture per acre.	Begin applications when birds begin feeding or crop begins to ripen. New York only: Aerial application is prohibited.

Migrate™

Specimen Label

USE SITE	APPLICATION DIRECTIONS	NOTES
Sunflowers	Apply by aerial application at a rate of 2.5 - 5 gallons of Rejex-4M Igate 0.1-0.2 lbs a.i./per acre or 7.5-15 fl. oz. per 1000 sq. ft. Use a 1:3 dilution with water and apply at rates of 10-20 gallons of spray mixture per acre.	Reapply 22 days after application. New York only: Aerial application is prohibited.

TURF AND HYDROSEEDING

**USE RESTRICTIONS:**  
Use this product to repel birds from golf courses, other turf areas (including around agricultural buildings and farmsteads), and hydroseeded areas. This product must be applied using appropriate spray equipment. Do not apply when grass is wet or rain is expected.

**DILUTION DIRECTIONS:**  
Shake or stir Rejex-4M Igate container well prior to diluting. For better coverage, add a superwetter (such as a non-fmt, organofunctional silicone surfactant, or sticker spreader), as directed on the label. Mix product outside or in a ventilated area. Mix Rejex-4M Igate with water at a ratio of 1 part product to 20 parts water. For example, mix 2 pints of product with 5 gallons of water to make 5.25 gallons of spray mixture.

**APPLICATION DIRECTIONS:**  
Apply at a rate of 1.2 gallons of spray mixture (0.7 lbs. a.i.) per 1000 sq. ft. of turf. Spray evenly on area to be protected to provide thorough coverage and allow mixture to dry before permitting human activity on the treated area. Repeat application in 4 days or as warranted by Canada goose activity. Do not mow treated area for several days after application.

GARBAGE, FOOD STUFFS, WASTE, AND TRASH

**USE RESTRICTIONS:**  
Use this product to repel birds from garbage, spilled grain, and other waste food stuffs. This product must be applied using appropriate spray equipment. Do not apply if rain is expected.

**DILUTION DIRECTIONS:**  
Shake or stir Rejex-4M Igate container well prior to diluting. For better coverage, add a superwetter (such as a non-fmt, organofunctional silicone surfactant, or sticker spreader), as directed on the label. Mix product outside or in a ventilated area. Mix Rejex-4M Igate with water at a ratio of 10-50% v/v with water. To make a 20% v/v mixture, mix 1 gallon of product with 5 gallons of water. To make a 50% v/v mixture, mix 1 gallon of product with 1 gallon of water.

**APPLICATION DIRECTIONS:**  
Spray mixture evenly on area to be protected to provide thorough coverage and allow mixture to dry before permitting human activity on the treated area. Repeat application as warranted by bird activity.

SPOT REPELLING

**USE RESTRICTIONS:**  
Use this product to immediately repel birds from iron structures, turf areas, landfills, and other outdoor areas (including around agricultural buildings and farmsteads). This product must be applied using appropriate spray equipment.

**DILUTION DIRECTIONS:**  
Shake or stir Rejex-4M Igate container well prior to diluting. Mix product outside or in a ventilated area. Mix Rejex-4M Igate with water at a ratio of 10-50% v/v with water. To make a 20% v/v mixture, mix 1 gallon of product with 5 gallons of water. To make a 50% v/v mixture, mix 1 gallon of product with 1 gallon of water.

**APPLICATION DIRECTIONS:**  
Spray mixture evenly on area to be protected to provide thorough coverage and allow mixture to dry before permitting human activity on the treated area. Repeat application as warranted by bird activity.

STORAGE & DISPOSAL
Do not contaminate water, food, or feed by storage or disposal. <b>PESTICIDE STORAGE:</b> Rejex-4M Igate should be stored in its original container in a cool, dry locked place out of reach of children and out of direct sunlight. Do not use or store near heat or open flame.
<b>PESTICIDE DISPOSAL:</b> To avoid waste, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (when such programs are run by state or local governments or by industry).
<b>CONTAINER DISPOSAL:</b> Non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Empty the remaining contents into application equipment or a mix tank and flush for 10 seconds, after the flow begins to drip. Fill the container 1/4 full with water and soap. Shake for 10 seconds. Pour mixture into application equipment or a mix tank or store mixture for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If burned, stay out of smoke.

WARRANTY STATEMENT

Notice: The statements on this label are believed to be true and accurate, but because conditions are beyond our control, Ceannad, Inc. does not make, nor does it authorize any agent or representative to make any oral, or written warranty, guarantee, or representation, express

or implied, concerning this material or the use thereof beyond the statements on the label. Ceannad, Inc. and its representatives hereby for, and expect the buyer to assume a risk for, any claim or personnel injury or property damage or loss resulting from handling, storage, or use of this material not in accordance with the written directions. No other warranty exists.