

# Microcide SQ™

## BROAD SPECTRUM DISINFECTANT

**HOSPITAL DISINFECTANT • CLEANS AS IT DISINFECTS**  
**CLEANER • DISINFECTANT • BACTERICIDE • VIRUCIDE\***  
**FUNGICIDE • DEODORIZER • MILDEWSTAT**

### ACTIVE INGREDIENTS:

Alkyl (60%C14, 30%C16, 5%C18) dimethyl benzyl ammonium chloride .....	2.37%
Alkyl (68%C12, 32%C14) dimethyl ethyl benzyl ammonium chloride .....	2.37%
INERT INGREDIENTS .....	95.26%
TOTAL .....	100.00%

**KEEP OUT OF REACH OF CHILDREN**  
**DANGER**

**SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT**



54 York Street Suite 1B • Portland, Maine 04107

1-800-755-8420

EPA Reg. No. 61178-1-70391 EPA Est. No. 70391

NET CONTENTS: 8 ounces

### PRECAUTIONARY STATEMENTS

## DANGER

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**KEEP OUT OF REACH OF CHILDREN:** Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothes. Harmful if swallowed. Wear goggles or face shield and rubber gloves when handling. Avoid contamination of food. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

**STATEMENT OF PRACTICAL TREATMENT:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eye contact, call a physician. If swallowed, drink egg whites, gelatin solution or if these are not available drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**DISINFECTION:** To be used to disinfect inanimate, hard, non-porous environmental surfaces such as walls, floors, sink tops, toilet bowls, tables, chairs, telephones, chair arms, bed frames, cabinets, bathtubs, and urinals.

**BOTTLE USE DILUTION:** Loosen cap on upper dilution chamber. Gently squeeze lower chamber until upper chamber is filled to 1/2 ounce level. Add 1/2 ounce of Microcide SQ™ per 32 ounces of water. **Bactericide:** Remove gross filth or heavy soil. Apply solution with a cloth, mop, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes and then let air dry. For heavily soiled areas, a pre-cleaning step is required. Microcide SQ™ is effective in hard water up to 400 ppm hardness (calculated as CaCO<sub>3</sub>) and in the presence of organic soil (5% blood serum). **Virucidal\*:** This product when used on inanimate, hard, non-porous, environmental surfaces at 1/2 ounce per 32 ounces of water for a 10 minute contact time (5% organic soil) is effective against Influenza A2-Asian Virus, HBV (Hepatitis B Virus), Herpes Simplex type 1 Virus, Herpes Simplex type 2 Virus, Pseudorabies Virus, and Vaccinia Virus. Kills HIV-1 (AIDS Virus) after 30 seconds contact time, and Polio virus type 1 (Chad strain) after 30 minute contact time (5% organic soil) **FOR CALIFORNIA USE ONLY:** HBV contact time is 15 minutes. \*Kills HIV-1 AND HBV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOODY/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV-1 (AIDS virus) or HBV (Hepatitis B Virus).

**MILDEWSTAT:** To control mold and mildew on pre-cleaned, hard, non-porous surfaces (such as floors, walls, table tops) add 1/2 ounce Microcide SQ™ per 32 ounces of water. Apply solution with a cloth, mop, or sponge making sure to wet all surfaces completely. Let air dry. Prepare fresh solution for each use. Repeat application at weekly intervals or when mildew growth reappears.

**DEODORIZER:** Microcide SQ™ deodorizes garbage storage areas, empty garbage bins and cans, toilet bowls and any other odor-causing areas. Use 1/2 oz. per 32 ounces of water.

**FUNGICIDE:** At 1/2 oz. per 32 ounces, Microcide SQ™ is effective against the pathogenic fungus Trichophyton mentagrophytes in 5% organic soil.

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 AND HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:** **PERSONAL PROTECTION:** Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks or eye coverings.

**CLEANING PROCEDURE:** Blood and other body fluids must be thoroughly cleaned from surfaces before application of the disinfectant.

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

**CONTACT TIME:** Leave surfaces wet for 30 seconds and 10 minutes for HIV-1 and HBV respectively. The 30-second contact time will not control other common types of viruses and bacteria. **FOR CALIFORNIA USE ONLY:** HBV contact time is 15 minutes.

**STORAGE AND DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Do not store on side. Avoid creasing or impacting of side walls.

**STORAGE:** Store in securely closed original container. Avoid storage at temperature extremes or in direct sunlight. Avoid shipping or storing below freezing. If product freezes, thaw at room temperature and shake gently to remix contents. Store in an area that will prevent cross contamination of other pesticides, fertilizer, food or feed.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse container. Replace cap and discard container in trash.