

Summer has arrived!



A COSTLY NUISANCE

Many birds molt their worn and damaged feathers in late summer. These discarded feathers can clog vents, drain spouts and rain gutters.

In summer, many species will feed on harvested crops, seeds and shrub-producing berries to become a costly nuisance to growers.

Summer is also the time when nesting birds raise their babies. Nests built in signage and near AC units cause damage.

During mid-to-late summer, many birds head for the wider countryside to feast on seasonal seeds and fruits.

RISKS IN LATE SUMMER

Woodpeckers can be particularly destructive to cedar and redwood siding in summer, especially to property that is vacant or unattended.

During late summer, crows gather from many miles to form communal night roosts that can damage property.

Blackbird summer roosts

can create a nuisance and health hazard to property owners.

Doves present risks in late summer when large roosting and loafing flocks gather for feeding, watering, and gritting on airport turf and runways, increasing the risk of bird-aircraft collisions



SOMETHING TO CROW ABOUT

Arriving in large, noisy numbers, crows often damage seedling corn plants by pulling the sprouts and consuming the

kernels. They also like to eat ripening grain sorghum, commercial sunflowers, pecans, various fruits, and watermelons.



SUMMER BREEDING SEASON

Swallows build mud nests inside buildings with open access barns, stables, sheds, carports and porches. These nests are unsightly and can damage paint.

During the summer breeding season, seagulls are drawn

to parking lots to feed on food scraps around supermarkets, restaurants, dumpsters, and garbage cans.



Keep summer birds off your property with Professional Grade Bird Control Products from **Bird B Gone**.

Made in the USA



BIRD•B•GONE[®]

Our customer support team is just a phone call away to assist you in finding the best solution for your budget.

Call 1-800-392-6915

birdbgone.com